



PNEUMONIA IN CHILDREN MODERN FEATURES

Nabiyeva F. S.

Senior Lecturer of the Department of Clinical Laboratory Diagnostics,

Muhitdinov O'. M.

Cadet of the Department of Clinical Laboratory Diagnostics,
Samarkand State Medical University

Abstract

Pneumonia in children remains a serious public health problem, requiring the introduction of new approaches to diagnostics. Modern research emphasizes the importance of a combination of clinical, laboratory and radiological methods. According to the World Health Organization, millions of cases occur each year, especially among infants and young children. The persistent high level of morbidity and mortality determines the relevance of the problem.

Keywords: Streptococcus pneumoniae, etiology, pathogenesis, laboratory diagnostics, instrumental diagnostics, prevention.

Pneumonia is one of the most common health problems among children and has serious consequences worldwide, especially in developing countries. This disease is an inflammation of the lungs caused by bacteria, viruses or pathogens. Pneumonia in children is predominantly infectious in nature, which further increases its prevalence.

Pneumonia caused by infectious agents is a dangerous disease of the lower respiratory tract that can be fatal.

The etiology of pneumonia in children is a complex and multifaceted process that includes many factors:

- Bacterial infections. The most common cause of pneumonia is bacteria. Bacteria such as Streptococcus pneumoniae, Haemophilus influenzae, and Mycoplasma pneumoniae can cause pneumonia in children. Bacterial infections usually develop quickly, and the inflammation can be severe.



- Viral infections. Viruses such as respiratory syncytial virus (RSV), influenza viruses, and adenoviruses can also cause pneumonia. Viral pneumonia is often less severe than bacterial pneumonia, but in some cases, especially in children with weakened immune systems, it can cause serious complications.
- Fungal infections. Fungi can also cause pneumonia, but this disease is more common in children with weakened immune systems, such as those with cancer or HIV infection.

Exposure to environmental factors such as air pollution, passive smoking and inadequate vaccination significantly increases the risk of developing pneumonia. Children's vulnerability to infections is explained by their immature immune system, which makes prevention and timely diagnosis especially important.

The pathogenesis of pneumonia in children is a multi-stage process involving various factors, ranging from the effects of pathogenic microorganisms to the individual characteristics of the child's immune system [2,8,15]. In younger children, whose immune systems are not yet fully developed, pneumonia can develop rapidly, requiring immediate intervention. The risk of developing severe forms of the disease increases in the presence of predisposing factors such as prematurity, chronic diseases or poor socioeconomic conditions.

Symptoms of pneumonia can vary depending on the age of the child, but the most common signs include fever, cough, shortness of breath, and difficulty breathing. The child may complain of chest pain, fatigue and general weakness. When pneumonia occurs, there is also an increase in heart rate and a change in skin color, which may take on a bluish tint. In some cases, children may experience vomiting and a reluctance to eat, which worsens their condition. It is important to note that symptoms may be less severe in infants, so parents should be aware of any changes in their baby's behavior, such as fussiness or refusal to feed [3,9,10].

Laboratory diagnostics of pneumonia in children is an important step in the process of early detection and effective treatment of this disease. The main goal of diagnostics is to determine the causative agent of the infectious process and assess the severity of the patient's condition. To begin with, a general blood test is performed, which helps to establish the presence of an inflammatory process, revealing leukocytosis and an increase in the level of C-reactive protein.



International Conference on Scientific Research in Natural and Social Sciences

Hosted online from New York, USA

Website: econfseries.com

2nd April, 2025

Microbiological testing, including sputum culture and nasopharyngeal swabs, helps identify bacterial agents, while PCR testing can provide information about viral or mycoplasma infections. In addition, blood gas testing provides information about the degree of respiratory failure. The combination of these laboratory diagnostic methods allows doctors not only to confirm the diagnosis of pneumonia, but also to choose the optimal therapeutic strategy for each child [5,14].

Instrumental diagnostics of pneumonia in children plays a key role in the timely and accurate detection of the disease. The most common method is chest X-ray, which allows visualization of inflammatory changes in the lungs. It helps differentiate pneumonia from other diseases such as bronchitis or pleurisy due to the clarity of the images showing areas of darkening or consolidation of the lung tissue. In addition, ultrasound examination of the lungs is actively used to assess the condition of the pleura and the presence of fluid in the pleural cavity. This method is safe and non-invasive, which makes it particularly attractive for children. Ultrasound can help diagnose complications of pneumonia, such as pleural effusion [6,11].

Computed tomography (CT) is used less frequently, but in complex cases it provides more detailed images and helps determine the extent of lung tissue damage. It is important to note that the choice of diagnostic method always depends on the clinical picture and condition of the patient, as well as the age of the child and the presence of concomitant diseases.

Prevention of pneumonia in children is an important task that requires a comprehensive approach. The main measures to prevent this disease are vaccination, compliance with hygiene standards and strengthening the immune system. Vaccination against pneumococcal infection and influenza helps reduce the risk of developing pneumonia, especially in young children who are more susceptible to infections. Strengthening the immune system also plays a key role. Proper nutrition, rich in vitamins and minerals, physical activity and hardening will help the body resist infections [11,13,14].



Conclusion

Thus, early diagnosis and prevention of pneumonia in children is not only relevant, but also necessary to reduce morbidity and improve the overall health of the younger generation.

REFERENCES:

1. Малахов А. Б. и др. Внебольничная пневмония у детей: алгоритмы диагностики и антибактериальной терапии //Практическая пульмонология. – 2019. – №. 2. – С. 18-26.
2. Сергеева Е. В., Петрова С. И. Внебольничная пневмония у детей. Современные особенности //Педиатр. – 2016. – Т. 7. – №. 3. – С. 5-10.
3. Бердиярова Ш.Ш., Юсупова Н.А. Особенности иммунометаболических нарушений иммунологической реактивности при гематогенных остеомиелитах. Вестник науки и образования, 29-32.
4. Клинико-лабораторная диагностика внебольничных пневмоний у детей ШШ Бердиярова, НА Юсупова, ХИ Ширинов Вестник науки и образования, 80-83.
5. Ибрагимов Б.Ф., Ибрагимова Н.С. Роль гомоцистеина в патогенезе синдрома поликистозных яичников у женщин International scientific review, Boston, USA. January 22-23, 2020.
6. Душанова Г.А., Набиева Ф.С., Садинова М.Ж., Нурматова Д.М. Анализ взаимосвязей параметров иммунного гомеостаза с состоянием системы ПОЛ-АОС // Вестник науки и образования, 2021. № 2 (105). Часть 2.
7. Nabiyeva F. S., Umirkulova S. I. DIAGNOSIS AND PREVENTION OF TORCH INFECTION IN PREGNANT WOMEN //Web of Medicine: Journal of Medicine, Practice and Nursing. – 2025. – Т. 3. – №. 1. – С. 76-78.
8. Umarova S.S., Nabiyeva F.S., Ibragimova N.S. Early diagnostics of echinococcosis in children // European research: innovation in science, education and technology. London, United Kingdom. January 9-10. С. 88-90, 2020.
9. Isomadinova L. K., Kudratova Z. E. Clinical and laboratory characteristics of vomiting in pregnant women in early pregnancy //Doctor's herald journal. – 2023. – Т. 2. - С. 52-56.



10. Kudratova Z. E., & Shamsiddinova M. Sh. (2023). LABORATORY METHODS FOR DIAGNOSING UROGENITAL CHLAMYDIA. Open Access Repository, 10 (10), 5–7.
11. Sabirovna I. N., Shekhrozovna B. F. DIAGNOSTIC CRITERIA AND TREATMENT OF TYPE 2 DIABETES MELLITUS //Galaxy International Interdisciplinary Research Journal. – 2023. – Т. 11. – №. 10. – С. 237-240.
12. Tursunov Feruz O'Ktam O'G'Li, Raximova Gulchiroy Olim Qizi, Isroilova Umidaxon, Turayeva Shaxnoza ASSESSMENT OF CARBOHYDRATE METABOLISM IN PATIENTS WITH DIABETES AND COVID-19 // ReFocus. 2022. №4.
13. Feruz O'ktam o'gli T., Mengdobilovich M. N. ANALYSIS OF GLYCEMIA AND GLUCOSURIA IN PATIENTS WITH DIABETES AND COVID-19 //Open Access Repository. – 2023. – Т. 4. – №. 2. – С. 177-181.
14. Набиева Ф. С., Ташанова З. И., Мухаммадиев Н. М. РАННЯЯ ДИАГНОСТИКА ИНФЕКЦИОННЫХ ЗАБОЛЕВАНИЙ НА СОВРЕМЕННОМ ЭТАПЕ //Бюллетень педагогов нового Узбекистана. – 2024. – Т. 2. – №. 1. – С. 8-11.
15. Nabieva F. S. et al. THE PROSPECT OF USING BAKERDREAM YEAST (SACCHAROMYCES CEREVISIAE) IN THE PRODUCTION OF ENZYMES.