



International Conference on Medical Science, Medicine and Public Health

Hosted online from Jakarta, Indonesia

Website: econfseries.com

30th October, 2025

THE EFFICIENCY OF DETECTION OF LATENT TUBERCULOSIS INFECTION IN CHILDREN

Rustamova Saodat Abdullaevna

Bukhara State Medical Institute, basic doctoral student of the Department of Phthisiology and Pulmonology

Muazzamov Bahodir Rakhmanovich

Bukhara State Medical Institute, Associate Professor of the Department of Phthisiology and Pulmonology, Candidate of Medical Sciences

Abstract

Considering the invaluable role of the body's immune system in the formation and development of this disease, it is important to determine and assess the immune status of children diagnosed with tuberculosis. Based on the above, the indicators of humoral immunity, which are among the indicators of the immune status of children with tuberculosis, were determined, and the results were interpreted and analyzed.

Keywords: tuberculosis, children, latent tuberculosis infection, diagnostics.

Relevance

The prevalence of tuberculosis among children is a variable indicator that characterizes the epidemiological situation with tuberculosis and depends on socio-economic, medical-biological factors, as well as the measures taken against tuberculosis in the region.

Latent tuberculosis infection (LTI) is a condition in which there is a LTI in the human body that causes positive reactions to immunological tests, including tuberculosis allergens, in the absence of clinical and radiological signs.

The purpose of the study: To increase the efficiency of detection of LTBI among children using immunological methods. To study the immunobiological state in children and adolescents.



E CONF SERIES



International Conference on Medical Science, Medicine and Public Health

Hosted online from Jakarta, Indonesia

Website: econfseries.com

30th October, 2025

Material and research methods. To solve the set tasks, 50 children with LTBI were analyzed; this contingent formed the main group of those observed. For comparative analysis, a control group was formed - 25 children who had contact with patients with tuberculosis and 25 children and adolescents with primary pulmonary tuberculosis who received treatment.

Both immunological and biochemical studies will be carried out in the laboratories of the above-mentioned clinics. A total of 100 children and adolescents will be examined.

Laboratory tests include: general blood and urine tests, biochemical blood tests. Determination of tuberculosis infection activity in children using immunological tests Diaskintest and QuantiFERON test. Bacterioscopic and bacteriological tests, X-ray examination (CT, MSCT of the lungs, as indicated), Immunobiological research, quantitative studies of procalcitonin, immunoglobulins and complement component C3. Quantitative and relative indicators of immunocompetent cells belonging to the T- and B-lymphocyte system, relative indicators of non-specific protective factors and quantitative indicators of humoral immunity parameters.

Results of the study. As a result of the conducted comprehensive examination including radiation methods, children from the main groups were divided into subgroups by the presence of a specific process, which allowed us to compare them. Comparison of subgroups of examined children from family and unspecified contact with various manifestations of tuberculosis infection. In children from family contact, chemoprophylaxis (CP) was carried out with the same frequency both in the A1 and A11 subgroups (35.0% (14) versus 36.7% (44)), however, children developed tuberculosis regardless of its implementation. At the same time, in group B, CP was prescribed to almost all children (82.3 (B1) and 88.9% (B11)). This fact may indirectly indicate the lack of alertness of phthisiologists for the development of the disease in the presence of contact, while an increase in sensitivity to tuberculin is a justification for prescribing preventive measures.

The analysis of concomitant pathology in subgroups A1 (healthy) and A11 (sick) children from the family focus of infection revealed no significant differences (67.5% and 58.3%, respectively). In group B, the number of children with



E CONF SERIES



International Conference on Medical Science, Medicine and Public Health

Hosted online from Jakarta, Indonesia

Website: econfseries.com

30th October, 2025

concomitant pathology was lower, but the prevalence of concomitant pathology in BP (48.1%) compared to B1 (32.1%) was also not found. A significantly high level of concomitant pathology was noted in sick patients from family contact (AII) compared to healthy children from an unspecified contact (B1) (58.3% (AII) versus 32.9% (B1), $X^2 = 12.33$ $p <0.001$).

The changes identified during the radiological examination require comparison with the immunological clinical methods used to determine the activity of tuberculosis infection in order to form a diagnosis.

Thus, the presence of concomitant pathology may be a risk factor for the development of tuberculosis in the presence of contact and does not have a significant impact on the development of tuberculosis in its absence.

LIST OF REFERENCES

1. Aslonov F. I., Rustamova S. A., Raxmonova K. M. Immunopathological aspects in patients with first detected pulmonary tuberculosis //World Bulletin of Public Health. – 2021. – T. 4. – C. 91-95.
2. Abdullayevna R. S. CLINICAL AND RADIOLOGICAL FEATURES OF NEWLY DETECTED PULMONARY TUBERCULOSIS IN PATIENTS WITH CONCOMITANT DISEASES //Intent Research Scientific Journal. – 2023. – T. 2. – №. 3. – C. 45-56.
3. Abdullaevna, R. S., & Rakhmanovich, M. B. (2023). Immunological Features of Pulmonary Tuberculosis in Patients with Drug Resistance. Scholastic: Journal of Natural and Medical Education, 2(4), 40-57.
4. Kh, Usmonov I., Bahodir R. Muazzamov, and Muhtor F. Jumaev. "Features of diagnostics and treatment of drug-resistant forms of pulmonary tuberculosis." International journal of pharmaceutical research 13.1 (2021): 2484-2489.